


Interactions and disease transmission between domestic cats and carnivores in Madagascar

1. How does dogs and cat occupancy probability and colonization rate affect Eupleridae occupancy probability
2. What are the variables associated *Toxoplasma gondii* exposure in Eupleridae



Encephalomyelitis by *Toxoplasma gondii* in a captive fossa (*Cryptoprocta ferox*)

J.M. Corpa^{a,*}, A. García-Quirós^a, M. Casares^b, A.C. Gerique^b, M.D. Carbonell^b, M.T. Gómez-Muñoz^c, F.A. Uzal^d, J. Ortega^a




Fig. 1. (a) Brain. Focus of inflammation composed by gliosis, vacuolated macrophages (Gitter cells), lymphocytes and plasma cells. Hematoxylin and eosin. Bar = 80 µm. (b) Brain. A *T. gondii* like tissue cyst. Hematoxylin and eosin. Bar = 20 µm. (c) Spinal cord. A *T. gondii* like tissue cyst in the white matter. Hematoxylin and eosin. Bar = 20 µm. (d) Brain. Protozoan cyst positive to anti-*Toxoplasma gondii* polyclonal antibody. Avidin-Biotin peroxidase complex (ABC). Bar = 20 µm.

Acknowledgments: Tanjona for mechanistic model, Mihaja for fieldwork

Mechanistic model: How does the occupancy of dogs & cats affect the occupancy of Eupleridae in forest patches.

States:

Eu: Eupleridae

Unoccupied

Eo: Eupleridae Occupied

Do: Dog occupied


Du: Dog unoccupied

Processes:



Dog colonization

Dog extinction

Eupleridae extinction



Statistical model: what are risk factors of exposure to *Toxoplasma gondii* in Eupleridae



- Response variable: pres/absence of antibodies to *Toxoplasma*
- Predictor variables: Species, Age, Sex, distance of capture to closest village, edge of forest
- Family: “Binomial”
- Link: logit
- Hypothesis: Exposure to *Toxoplasma gondii* in Eupleridae is associated with demographic and spatial factors in Eupleridae of Madagascar

Variable	P-value	Odds ratio	95% CI
Sex	0.03		
Female		1.00	
Male		4.13	1.16–14.78
Species	0.003		
<i>G. fasciata</i>		1.00	
<i>Cr. ferax</i>		24.55	2.19–275.47
<i>G. elegans</i>		5.07	1.31–19.58

```
toxofull<-glm(Toxo ~Species+Sex+distance_vill+distance_edge, family = binomial, data = euplerid)
```

Patterns of Exposure of Carnivores to Selected Pathogens in the Betampona Natural Reserve Landscape, Madagascar

Fidisoa Rasambainarivo,^{1,2,6} Mamy Navalona Andriamihajarivo,³ Edward Dubovi,⁴ and Patricia G Parker^{1,5} ¹Department of Biology and Harris World Ecology Center, University of Missouri Saint-Louis, One University

Next Steps

1. Collect samples from additional sites in multiple regions of Madagascar.
2. Identify other pathogens that may affect the populations of Eupleridae in protected areas.

